

BIJLAGE 1

LITERATUURLIJST



LITERATUURLIJST

Hoofdstuk 1-4

Agreement on the Conservation of Small Cetaceans in the Baltic, North East Atlantic, Irish and North Seas).

Convention on the Conservation of Migratory Species of Wild Animals

Agreement on the Conservation of Populations of European Bats

Agreement for the Conservation of Seals in the Wadden Sea

Desk study, Helicopter accessibility of oil & gas platforms near the offshore wind farms Hollandse Kust (zuid and noord), april 2017

ECN, 2018, Optimal wind farm power density analysis for future offshore wind farms

ECN-E--18-025, Study into levelised Cost of Energy of variants for wind farm site boundaries of Hollandse Kust (west), Ten Noorden van de Waddeneilanden and IJmuiden

Hintzen et al, in voorbereiding.

Kader Ecologie en Cumulatie 3.0

Kamerstukken II, 2013/14, 33 450, nr. 22

Kamerstukken I/II, 2014/15, 33 561, A/nr. 11 (herdruk).

Kamerstukken II, 2015/16, 31 710, nr. 45

Kamerstukken II, 2016/17, 31 510, nr. 64

Kamerstukken II, 2017/18, 33 561, nr. 42

Ministerie van Infrastructuur en Milieu, Beleidsnota Noordzee 2016-2021, 2015

Ministerie van Infrastructuur en Milieu (2011), Integraal Beheerplan Noordzee 2015.

Ministerie van Infrastructuur en Milieu, Nationaal Waterplan (2016-2021), NWP2, 2015

Ministerie van Infrastructuur en Milieu, Riksstructuurvisie Windenergie op Zee, partiële herziening van het NWP1, 26 september 2014 (Kamerstukken I/II, 2014/15, 33 561, A/nr. 11 (herdruk) Kamerstukken II 2013/14, 31 510, nr. 49.

Ministerie van Infrastructuur en Milieu, Rijksstructuurvisie Windenergie op Zee aanvulling gebied Hollandse Kust, 2016

Ministerie van V&W, VROM en LNV, Beleidsnota Noordzee, 2009.

Ministerie van V&W, VROM en LNV (2009), Nationaal Waterplan 2009-2015, NWP1, 22 december.

NLR-CR-2016-266 Offshore windturbinezog en veilige helikopteroperaties

OSPAR Commission (2008) OSPAR List of Threatened and/or Declining Species and Habitats. Reference Number: 2008-6.

RICHTLIJN 2014/89/EU Richtlijn Maritieme Ruimtelijke Ordening

RvO, 2018, Geological Desk Study Hollandse Kust (west) Wind Farm Zone, WOZ2180087.

SER, Energieakkoord voor duurzame groei, 2013.

Vanossmael, C., K.A. Willems, D. Claeys, M. Vincx & C. Heip 1982. Macrobenthos of a sublittoral sandbank in the South-ern Bight of the North Sea. J. mar. biol. Ass. U.K. 62: 521-534

Hoofdstuk 5 Morfologie en hydrologie

ACRB. (2013). Mogelijke morfologische effecten van het Prinses Amaliawindpark. Rapport Q214R1

Carpenter J.R. et al. (2016) Potential impacts of offshore wind farms on North Sea stratification, PLoS one, 11(8): p. e0160830.

Chakrabarti, S.K. (1987) Hydronamics of offshore structures, WitPress, Billerica USA

Danish Hydraulic Institute (1999) Horns Rev Wind Power Plant, Environmental impact assessment of hydrography.

Dankers, P.J.T. (2015) Notitie slibverspreiding loswallen. Report No.: RDCHM_BE1044_N0003_902199_f02

Van Dijk (2018), Gaida et al. (2018): Timo C. Gaida, Mirjam Snellen, Thaiënne A. G. P. van Dijk, Dick G. Simons 2018Geostatistical modelling of multibeam backscatter for full-coverage seabed sediment maps. (<http://creativecommons.org/licenses/by/4.0>)

Deltares (2018A), Raaijmakers, T. ,Roetert, T., Bruinsma, N. Riezebos h.J., Van Dijk, H, Forzoni A, Vergauwen S, Grasmeijer B., 2018. Morphodynamics and scour mitigation for Hollandse Kust (noord) Wind Farm Zone. Report nr: 11202796-000

Deltares (2018B): Tom Roetert, Niek Bruinsma, Chris Mesdag, 2018; Seabed mobility assessment Hollandse Kust (west-alpha), report nr: 11202582-002

Floeter, J., van Beusekom, J.E.E., Auch, D., Callies, U., Carpenter, J., Dudeck, T., Eberle, S., Eckhardt, A., Gloe, D., Hänselmann, K., Hufnagl, M., Janßen, S., Lenhart, H., Möller, K.O., North, R.P., Pohlmann, T., Riethmüller, R., Schulz, S., Spreizenbarth, S., Temming, A., Walter, B., Zielinski, O., Möllmann, C. (2017). Pelagic effects of offshore wind farm foundations in the stratified North Sea. Progress in Oceanography. 156, pp. 154-173.

Hoffman, C.J.C.M. en H.J. Verheij, Scour Manual, A.A. Balkema Publications Korevaar, 1990

De Kok (2004) Slibtransport langs de Nederlandse Kust. Bronnen, fluxen en concentraties. RIKZ/OS/2004.148w

NationaleVisieKust (2013) Deltaprogramma Kust, Nationale visie kust, Den Haag

Niessen, A.C.H.M. en Schüttenhelm, R.T.L, 1986 - Oppervlakteelfstoffen/Raw materials at or near the surface. Rijks Geologische Dienst, Haarlem, map.

Van Rijn. L, 2015. PRINCIPLES OF SEDIMENTATION AND EROSION ENGINEERING IN RIVERS, ESTUARIES AND COASTAL SEAS (update 2015). Aqua publications.

Royal Haskoning (2005) EIS, SEA and Habitat Assessment for BritNed Interconnector

Salden, R.M., 1998. Een model voor het transport van slib in de Nederlandse kustzone. Een hulpmiddel naar scenario studies naar kustuitbreidingsplannen. Rapportage van SILTMAN deelprojecten 11 & 14. OS-98.119X, RIKZ

Sediment Atlas, RIKZ (2002) Ecologische Atlas Waddenzee

Svasek (2019): S.E. Poortman, M.J.G. van den Boomgaard, 2019. Seabed mobility study update cable route Windpark Hollandse Kust (noord). report 1976/U19159/B/SPo

Tennet. 2015. 66 kV Systems for Offshore Wind Farms. Report No.: 113799-UKBR-R02, Rev 2. 05-03-2015

Weerst, A.H. & F.L.M. Diermanse. Golfstatistiek op relatief diep water 1979-2002. WL|Delft. Rapport Q3770, december 2004

Wijnberg, K.M. (1995) Morphological behaviour of a barred coast over a period of decades. Ph. D. thesis Utrecht University.

Hoofdstuk 6 Vogels (zie ook de literatuurlijst die bij bijlage 4 hoort)

Arts, F.A. 2013. Trends en verspreiding van zeevogels en zeezoogdieren op het Nederlands Continentaal Plat 1991 – 2012. RWS-Centrale Informatievoorziening BM 13.28, Lelystad.

Barclay, R.M.R., E.F. Baerwald & J.C. Gruver, 2007. Variation in bat and bird fatalities at wind energy facilities: assessing the effects of rotor size and tower height. Canadian Journal of Zoology-Revue Canadienne De Zoologie 85(3): 381-387.

Baerwald, E.F., G.H. D'Amours, B.J. Klug & R.M.R. Barclay, 2008. Barotrauma is a significant cause of bat fatalities at wind turbines. Current Biology 18: R695-696.

Bearwald E.F., Edworthy J., Holder M., and Barclay R.M.R. 2009. A large scale mitigation experiment to reduce bat fatalities at wind energy facilities. J. Wildlife Management 73: 1077-1081.

Bat Conservation Trust 2017. http://www.bats.org.uk/pages/national_nathusius_pipistrelle_project.html .

Boshamer, J.P.C., & J.P. Bekker, 2008. Natusius' pipistrelles (*Pipistrellus natusius*) and other species of bats on offshore platforms in the Dutch sector of the North Sea. Lutra 51: 17-36.

Boele, A., J. van Bruggen, F. Hustings, K. Koffijberg, J.W. Vergeer & C. Plate, 2014. Broedvogels in Nederland in 2012. Sovon-rapport 2014/13. Sovon Vogelonderzoek Nederland, Nijmegen.

Boele, A., J. van Bruggen, F. Hustings, K. Koffijberg, J.W. Vergeer & T. van der Meij, 2016. Broedvogels in Nederland in 2014. Sovon-rapport 2016/04. Sovon Vogelonderzoek Nederland, Nijmegen.

Boele, A., J. van Bruggen, F. Hustings, K. Koffijberg, J.W. Vergeer & T. van der Meij, 2017. Broedvogels in Nederland in 2015. Sovon-rapport 2017/04. Sovon Vogelonderzoek Nederland, Nijmegen.

Boonman, M. 2018. Mitigerende maatregelen voor vleermuizen in offshore windparken. Evaluatie en verbetering van stilstandvoorziening. Bureau Waardenburg Raporthr. 18-278. Bureau Waardenburg, Culemborg.

Bouma S., & W. Lengkeek 2009. Development of underwater flora- and fauna communities on hard substrates of the offshore wind farm Egmond aan Zee (OWEZ). Report OWEZ_R_266_T1_20090126.

Bouma S. & W. Lengkeek 2012. Benthic communities on hard substrates of the offshore wind farm Egmond aan Zee (OWEZ). Including results of samples collected in scour holes. Report OWEZ_R_266_T1_20120206_hard_substrate.

- Bowgen, K. & A. Cook, 2018. Bird Collision Avoidance: Empirical evidence and impact assessments. JNCC Report No: 614. JNCC, Peterborough.
- Bradbury, G., Trinder, M., Furness, B., Banks, A.N., Caldow, R.W.G., Hume, D. 2014. Mapping Seabird Sensitivity to Offshore Wind Farms. PLoS ONE 9(9): e106366. doi:10.1371/journal.pone.0106366
- Bruynzeel, L.W., J. van Belle, L. Davids m.m.v F. van de Laar 2009. The impact of conventional illumination of offshore platforms in the North Sea on migratory bird populations. A&W report 1227, Altenburg & Wymenga Ecological Consultants, Veenwouden, 38 p.
- Camphuysen C.J., 2013. A historical ecology of two closely related gull species (Laridae): multiple adaptations to a man - made environment. Ph.D.-thesis, Univ. Groningen, Groningen.
- Camphuysen C.J. & A. Gronert, 2012. Apparent survival and fecundity of sympatric Lesser Black-backed Gulls and Herring Gulls with contrasting population trends. Ardea 100: 113-122.
- Collier, M.P., S. Dirksen & K.L. Krijgsveld, 2011. A review of methods to monitor collisions or micro-avoidance of birds with offshore wind turbines. Part 1: Review. Strategic Ornithological Support Services Project SOSS-03A. Rapport 11-078. Bureau Waardenburg, Culemborg.
- Collier, M.P., S. Dirksen & K.L. Krijgsveld, 2012. A review of methods to monitor collisions or micro-avoidance of birds with offshore wind turbines: Part 2: Feasibility study of systems to monitor collisions. Strategic Ornithological Support Services Project SOSS-03A. Rapport 11-215. Bureau Waardenburg, Culemborg.
- Cryan P.M. & R.M.R. Barclay, 2009. Causes of bat fatalities at wind turbines: hypotheses and predictions. Journal of Mammalogy 90: 1330–1340.
- Cryan, P.M., P. M. Gorresen, C. D. Hein, M. R. Schirmacher, R.H. Diehl, M.M. Huso, D.T.S. Hayman, P.D. Fricker, F.J. Bonaccorso, D.H. Johnson, K. Heist & D.C. Dalton, 2014. Behavior of bats at wind turbines. PNAS 111(42): 15126-15131.
- Cryan P.M. & R.M.R. Barclay, 2009. Causes of bat fatalities at wind turbines: hypotheses and predictions. Journal of Mammalogy 90: 1330–1340.
- Drewitt, A.L. & R.H.W. Langston, 2006. Assessing the impacts of wind farms on birds. Ibis 148: 29-42.
- Dirksen, S. 2017. Review of methods and techniques for field validation of collision rates and avoidance amongst birds and bats at offshore wind turbines. Sjoerd Dirksen Ecology / Rijkswaterstaat WVL.
- Dürr, T., 2013. Fledermausverluste an Windenergieanlagen. Daten aus der zentralen Fundkartei der Staatlichen Vogelschutzwarte im Landesumweltamt Brandenburg. Stand 25.09.2013. www.mluv.brandenburg.de/cms/media.php/.../wka_fmaus.xls.

Everaert, J. & E. Stienen, 2007. Impact of wind turbines on birds in Zeebrugge (Belgium). Significant effect on breeding tern colony due to collisions. *Biodiversity and Conservation* 16: 3345-3359.

Exo, K.M., O. Hüppop & S. Garthe. 2002. Offshore-Windenergieanlagen und Vogelschutz, Seevögel, Zeitschr, Verein Jordsand, Hamburg. 23: 83-95.

Fijn, R.C., K.L. Krijgsveld, M.J.M. Poot & S. Dirksen, 2015. Bird fluxes at risk altitudes in a Dutch offshore wind farm continuously measured with vertical radar. *Ibis* 157: 558-566.

Fijn, R.C., A. Gyimesi, M.P. Collier, J.C. Kleyheeg-Hartman, M. Boonman, J.W. de Jong & M.J.M. Poot, 2015b. Achtergronddocument ten behoeve van MER en PB windenergiegebied Borssele. Kavel I en II: vogels en vleermuizen. Rapportnr. 14-263. Bureau Waardenburg, Culemborg.

Fijn, R.C., A. Gyimesi, J.C. Kleyheeg-Hartman, M. Boonman & J.W. de Jong, 2015c. Achtergronddocument ten behoeve van MER en PB windenergiegebied Borssele. Kavel III en IV: vogels en vleermuizen. Rapportnr. 14-263. Bureau Waardenburg, Culemborg.

Gyimesi, A. & R. Lensink, 2012. Non-breeding adults or 'floaters' in bird populations. Rapport 11-200. Bureau Waardenburg, Culemborg.

Gyimesi, A. & R.C. Fijn, 2015a. Slachtofferberekeningen voor een windpark met een totaalvermogen van 380 MW in Kavel I of II in windenergiegebied Borssele. Bureau Waardenburg notitie 15.03297.

Gyimesi, A. & R.C. Fijn, 2015b. Slachtofferberekeningen voor windparken in de zuidelijke Noordzee met bestaande en geplande turbinetypes. Bureau Waardenburg notitie 15.03620.

Gyimesi, A. & R.C. Fijn, 2015c. Slachtofferberekeningen voor drie gebiedsvarianten van de uitbreiding van windenergiegebied Hollandse Kust. Bureau Waardenburg notitie 15.05577.

Gyimesi, A., M. Dorenbosch, J.W. de Jong, M. Boonman M. Teunis & R.C. Fijn, 2016. Achtergronddocument ten behoeve van MER en PB windenergiegebied Hollandse Kust. Kavel I en II: vogels, vleermuizen, vissen en benthos. Rapport 16-007. Bureau Waardenburg, Culemborg.

Gyimesi, A., J.W. de Jong, M.P. Collier, W. Bouten & R.C. Fijn, 2017a. Validation of biological variables for use in the SOSS Band model for Lesser Black-backed Gull *Larus fuscus* and Herring Gull *Larus argentatus*. Rapport 16-042. Bureau Waardenburg, Culemborg.

Gyimesi, A., T.J. Evans, J.F. Linnebjerg, J.W. de Jong, M.P. Collier & R.C. Fijn, 2017b. Review and analysis of tracking data to delineate flight characteristics and migration routes of birds over the Southern North Sea. Rapport 16-139. Bureau Waardenburg, Culemborg.

Gyimesi, A., J.L. Leemans, 2018. Adding OWEZ and PAWP to the KEC 3.0 calculations. MEMORANDUM, Bureau Waardenburg, Culemborg.

Gyimesi, A., R.P. Middelveld, B.M.C. Grutters, E. Stienen & R.C. Fijn, 2018a. Effects of Offshore Wind farms on the Behaviour of Lesser Black-backed Gulls. Report 17-175. Bureau Waardenburg, Culemborg.

Gyimesi, A., J.W. de Jong, A. Potiek & E.L. Bravo Rebolledo, 2018b. Actualisatie van KEC vogelaanvaring berekeningen volgens Routekaart 2030. Rapportnr. 18-290. Bureau Waardenburg, Culemborg.

Gyimesi, A., E.L. Bravo Rebolledo, J.C. Kleyheeg-Hartman, J.W. de Jong, M. Teunis, K. Didderen, M. Boonman, M. Schutter & R.C. Fijn, 2018c. Achtergronddocument ten behoeve van MER en PB windenergiegebied Hollandse Kust (noord). Kavel V en kavel VI: vogels, vleermuizen, vissen en benthos. Culemborg, Bureau Waardenburg.

Haan, D. de, D. Burggraaf, S. Ybema & R. HilleRisLambers 2007. Underwater sound emissions and effects of the pile driving of the OWEZ windfarm facility near Egmond aan Zee (Tconstruct). Report OWEZ_R_251_TC 20071029, Imares Wageningen UR.

Hal, R. van, B. Couperus, S. Fassler, S. Gastauer, B. Griffioen, N. Hintzen, L. Teal, O. van Keeken & E. Winter 2012. Monitoring- and Evaluation Program Near Shore Wind farm (MEPN SW). Fish community. IMARES Report C059/12. OWEZ_R_264_T1_20121215_final_report_fish. Imeres Wageningen UR.

Heinänen, S. & H. Skov, 2018. Offshore Wind Farm Eneco Luchterduinen Ecological Monitoring of Seabirds T3 (Final) Report. ENECO.

Hill, R., K. Hill, R. Aumuller, A. Schulz, T. Dittman, C. Kulekmeyer & T. Coppack 2014. Of birds, blades an barriers: Detecting and analysing mass migration events at alpha ventus. In: BSH & BMU 2014. Ecological Research at the Offshore Windfarm alpha ventus. Challenges, Results and Perspectives. Federal Maritime and Hydrographic Agency (BSH), Federal Ministry for the Environment, Nature Conservation and Nuclear Safety (BMU). Springer Spektrum.

Jain, A.A., R.R. Koford, A.W. Hancock & G.G. Zenner, 2011. Bat mortality and activity at a northern Iowa wind resource area. American Midland Naturalist 165: 185-200.

Jansen, H.W. & C.A.F. de Jong 2014. Underwater noise measurements in the North Sea in and near the Princess Amalia Wind Farm in operation. TNO report - TNO 2013 R11916. Den Haag, the Netherlands.

Johnston, A., A.S.C.P. Cook, L.J. Wright, E.M. Humphreys & N.H.K. Burton 2014, Modelling flight heights of marine birds to more accurately assess collision risk with offshore wind turbines. Journal of Applied Ecology 51: 31–41. doi: 10.1111/1365-2664.12191

Jonge Poerink, B., Lagerveld, S. & H. Verdaat, 2013. Pilot study Bat activity in the Dutch offshore wind farm OWEZ and PAWP. IMARES report number C026/13. The Fieldwork Company report number 20120402. IMARES Wageningen UR.

Krijgsveld, K.L., R.C. Fijn & R. Lensink, 2015. Occurrence of peaks in songbird migration at rotor heights of offshore wind farms in the southern North Sea. Report nr 15-119. Bureau Waardenburg, Culemborg

Krijgsveld K.L. 2014. Avoidance behaviour of birds around offshore wind farms. Overview of knowledge including effects of configuration. Report Bureau Waardenburg 13-268, 30p., Bureau Waardenburg, Culemborg, Netherlands.

Krijgsveld K.L., R.C. Fijn, M. Japink, P.W. van Horssen, C. Heunks, M.P. Collier, M.J.M. Poot, D.Beuker & S. Dirksen 2011. Effect studies Offshore Wind Farm Egmond aan Zee: Final report on fluxes, flight altitudes and behaviour of flying birds. NoordzeeWind report nr OWEZ_R_231_T1_20111114_flux&flight. Report nr 10-219. Bureau Waardenburg, Culemborg.

Lagerveld S., B .Jonge Poerink, & H. Verdaat 2014a. Monitoring bat activity in offshore wind farms OWEZ and PAWP in 2013. IMARES Report C165/14.

Lagerveld S., B. Jonge Poerink, R. Haselager & H. Verdaat 2014b. Bats in Dutch offshore wind farms in autumn 2012. Lutra 57 (2): 61-69.

Lagerveld S., B. Jonge Poerink, P. de Vries & M. Scholl, 2015. Bat activity at offshore wind farms LUD and PAWP in 2015. IMARES Report C001/2016.

Lagerveld, S., Janssen, R., Manshanden, J., Haarsma, A-J., de Vries, S., Brabant R., & Scholl, M. 2017a. Telemetry for migratory bats – a feasibility study; Wageningen, Wageningen Marine Research (University & Research Centre), Wageningen Marine Research report C011/17.

Lagerveld, S., H.J.G.A. Limpens, M.J. Schillemans & M. Scholl 2017b. Bat 1: Estimate of bat populations at the southern North Sea. Supporting note to ZDV report no. 2016.031 Migrating bats at the southern North Sea. Wageningen, Wageningen Marine Research (University & Research Centre), Wageningen Marine Research report no. C014.17/Dutch Mammal Society report no. 2017.08. 14 pp.

Lagerveld S., D. Gerla, J.T. van der Wal, P. de Vries, R. Brabant, E. Stienen, K. Deneudt, J. Manshanden & M. Scholl, 2017c. Spatial and temporal occurrence of bats in the southern North Sea area. Wageningen Marine Research (University & Research centre), Wageningen Marine Research report C090/17; 52 p.

Larsen, J.K. & M. Guillemette, 2007. Effects of wind turbines on flight behaviour of wintering common eiders: implications for habitat use and collision risk. Journal of Applied Ecology 44: 516-522.

Larsen, J.K. & M. Guillemette, 2007. Effects of wind turbines on flight behaviour of wintering common eiders: implications for habitat use and collision risk. Journal of Applied Ecology 44: 516-522.

Lensink R. & J. van der Winden 1997. Trek van niet-zeevogels langs en over de Noordzee: een verkenning. Rapport 97-023, Bureau Waardenburg, Culemborg.

Lensink, R. & P.W. van Horssen, 2012. Een matrixmodel om effecten op een populatie te voorspellen van slachtoffers door windturbines. Rapport 11-198. Bureau Waardenburg, Culemborg.

Lensink, R., J.W. de Jong & J.C. Kleyheeg-Hartman, 2015. Luchtfoto inventarisatie van op daken broedende meeuwen in Den Haag in 2010. Limosa 88: 114-124.

Leopold, M.F. & C.J. Camphuysen. 2007. Did the pile driving during the construction of the Offshore Wind Farm Egmond aan Zee, the Netherlands, impact local seabirds. Rapport Wageningen. IMARES Nr. C062/07 - NoordzeeWind Rapport OWEZ_R_221_Tc_20070525 aan Nuon Energy Sourcing.

Leopold M.F., E.M. Dijkman, L. Teal & the OWEZ-team 2010. Local birds in and around the Offshore Wind Farm Egmond aan Zee (OWEZ). NoordzeeWind rapport OWEZ_R_221_T1_20100731_local_birds. Imares / NoordzeeWind, Wageningen / IJmuiden.

Leopold, M.F., R. van Bemmelen, A. Zuur. 2012. Responses of Local Birds to the Offshore Wind Farms PAWP and OWEZ off the Dutch mainland coast. IMARES Report nr. C151/12. IMARES, IJmuiden.

Leopold, M.F., M.P. Collier, A. Gyimesi, R. Jongbloed, M.J.M. Poot, J. Tjalling van der Wal & M. Scholl, 2015. Iteration cycle: Dealing with peaks in counts of birds following active fishing vessels when assessing cumulative effects of offshore wind farms and other human activities in the Southern North Sea. Additional note to IMARES report number C166/14. Imares Wageningen UR / Bureau Waardenburg Culemborg.

Leopold, M.F., E.M. Dijkman, T. Teal & the OWEZ-team, 2011. Local Birds in and around the Offshore Wind Farm Egmond aan Zee (OWEZ) (T-0 & T-1, 2002-2010). IMARES Report nr. C187/11, NoordzeeWind report OWEZ_R_221_T1_20111220_local_birds. IMARES, IJmuiden.

Leopold, M.F., R. van Bemmelen & A. Zuur, 2013b. Responses of Local Birds to the Offshore Wind Farms PAWP and OWEZ off the Dutch mainland coast. IMARES Report nr. C151/12. IMARES, IJmuiden.

Leopold, M.F., M. Booman, M.P. Collier, N. Davaasuren, R.C. Fijn, A. Gyimesi, J. de Jong, R. Jongbloed, B. Jonge Poerink, J.C. Kleyheeg-Hartman, K.L. Krijgsveld, S. Lagerveld, R. Lensink, M.J.M. Poot, J. Tjalling van der Wal & M. Scholl. 2015. A first approach to deal with cumulative effects on birds and bats of offshore wind farms and other human activities in the Southern North Sea. Report number C166/14. Imares Wageningen UR, Wageningen.

Limpens, H.J.G.A., M. Boonman, F. Korner-Nievergelt, E.A. Jansen, M. van der Valk, M.J.J. La Haye, S. Dirksen & S.J. Vreugdenhil, 2013. Wind turbines and bats in the Netherlands – Measuring and predicting. Report 2013.12, Zoogdiervereniging & Bureau Waardenburg.

Limpens, H.J.G.A., S. Lagerveld, I. Ahlén, D. Anzionnat, T. Aughney, H.J. Baagøe, , L. Bach, P. Bach, J.P.C. Boshamer, K. Boughey, T. Le Campion, M. Christensen, J.J.A. Dekker, T. Douma,

M.-J. Dubourg-Savage, J. Durinck, M. Elmeros, A.-J. Haarsma, J. Haddow, D. Hargreaves, J. Hurst, E.A. Jansen, T.W. Johansen, J. de Jong, D. Jouan, J. van der Kooij, E.-M. Kyheroinen, F. Mathews T.C. Michaelsen, J.D. Møller, G. Petersons, N. Roche, L. Rodrigues , J. Russ, Q. Smits, S. Swift, E.T. Fjederholt, P. Twisk, B. Vandendriesche & M.J. Schillemans, 2017. Migrating bats at the southern North Sea - Approach to an estimation of migration populations of bats at southern North Sea . Rapport 2016.031. Zoogdiervereniging, Nijmegen/ Wageningen Marine Research.

Lindeboom, H.J., Kouwenhoven, H.J., Bergman, M.J.N, Bouma, S., Brasseur, S., Daan, R., Fijn, R.C., de Haan, D., Dirksen, S., van Hal, R., Hille Ris Lambers, R., ter Hofstede, R., Krijgsveld, K.L., Leopold M. & Scheidat M. 2011. Short-term ecological effects of an offshore wind farm in the Dutch coastal zone; a compilation. Environmental Research Letters 6 035101 doi: 10.1088/1748-9326/6/3/035101.

LWVT/SOVON 2002. Vogeltrek over Nederland 1976-1993. Schuyt & Co, Haarlem.

Maclean, I.M.D., Wright, L.J., Showler, D.A. & M.M. Rehfisch, 2009. A review of assessment methodologies for offshore windfarms. BTO Report commissioned by COWRIE Ltd. Marquenie, J., van der Laar, F., Poot, H. 2009. Groen Licht voor Vogels. De Levende Natuur 110(6): 290-291.

Marquenie, J., van der Laar, F., Poot, H. 2009. Groen Licht voor Vogels. De Levende Natuur 110(6): 290-291.

Masden, E. A., Haydon, D. T., Fox, A. D., Furness, R. W., Bullman, R., and Desholm, M. 2009. Barriers to movement: impacts of wind farms on migrating birds. – ICES Journal of Marine Science, 66: 746–753.

Masden, E.A., D.T. Haydon, A.D. Fox & R.W. Furness, 2010. Barriers to movement: Modelling energetic costs of avoiding marine wind farms amongst breeding seabirds. Mar Pollut Bull 60(7): 1085-1091.

Masden, E.A., Reeve, R., Desholm, M., Fox, A. D., Furness, R.W., Haydon, D.T. 2012. Assessing the impact of marine wind farms on birds through movement modelling. Journal of The Royal Society Interface 9: 2120-2130.

May, R. O. Reitan, K. Bevanger, S.-H. Lorentsen & T. Nygård 2015. Mitigating wind-turbine induced avian mortality: Sensory, aerodynamic and cognitive constraints and options. Renewable and Sustainable Energy Reviews 42: 170-181.

Niermann, I., R. Brinkmann, F. Korner-Nievergelt & O. Behr, 2011. Systematische Schlagopfersuche - Methodische Rahmenbedingungen, statistische Analyseverfahren und Ergebnisse. In Brinkmann et al. 2011.

Nuijten, R.J.M., K.A. Wood, T. Haitjema, E.C. Rees & B.A. Nolet, 2018. "Phenological changes in a migratory swan coping with climate change." 6th International Swan Symposium. Abstracts 52.

Nuijten, R.J.M., K.A. Wood, T. Haitjema, E.C. Rees & B.A. Nolet, 2018b. "Migratory swans adapting to climate change: short-stopping or short-staying?" 6th International Swan Symposium. Abstracts 69.

Petersen I.K. & Fox A.D. 2007. Changes in bird habitat utilisation around the Horns Rev 1 offshore wind farm, with particular reference on Common Scoter. NERI Report. National Environmental Research Institute, Ministry of the Environment, Rønde, Denmark.

Petersen, I.K., T.K. Christensen, J. Kahlert, M. Desholm & A.D. Fox, 2006. Final results of bird studies at the offshore wind farms at Nysted and Horns Rev, Denmark. NERI Report National Environmental Research Institute, Ministry of the Environment, Rønde, Denmark.

Plonczkier P. & I.C. Simms 2012. Radar monitoring of migrating pink-footed geese: behavioural responses to offshore wind farm development. *Journal of Applied Ecology* 49: 1187-1194. doi: 10.1111/j.1365-2664.2012.02181.x.

Poot, H., B.J. Ens, H. de Vries, M.A.H. Donners, M.R. Wernand & J.M. Marquenie 2008. Green light for nocturnally migrating birds. *Ecology and Society* 13(2): 47.

Poot M.J.M., R.C. Fijn R.J. Jonkvorst C. Heunks M.P. Collier, J. de Jong & P.W. van Horssen, 2011a. Aerial surveys of seabirds and marine mammals in the Dutch North Sea May 2010 – April 2011. Distribution in relation to future offshore wind farms. Report 10-235. Bureau Waardenburg, Culemborg.

Poot M.J.M., P.W. van Horssen, M.P. Collier, R. Lensink & S. Dirksen, 2011b. Effect studies Offshore Wind Egmond aan Zee: cumulative effects on seabirds. A modelling approach to estimate effects on population levels in seabirds. Report 11-026, Bureau Waardenburg, Culemborg.

Poot M.J.M., 2013. Eerste gedachten over de haalbaarheid van nieuwe nearshore windparken in Nederland in relatie tot mogelijke effecten op vogels. Notitie 13.01476 in opdracht van Pondera Consult. Bureau Waardenburg, Culemborg.

Rebke, M., V. Dierschke, C.N. Weiner, R. Aumüller, K. Hill & R. Hill, 2019. Attraction of nocturnally migrating birds to artificial light: The influence of colour, intensity and blinking mode under different cloud cover conditions. *Biological Conservation*, 233: 220-227.

Reid, J.B. & C.J. Camphuysen 1998. The European Seabirds at Sea database. Biol. Cons.

Rijkswaterstaat 2015. Kader Ecologie en Cumulatie t.b.v. uitrol windenergie op zee Deelrapport B - Bijlage Imares onderzoek Cumulatieve effecten op vogels en vleermuizen. Ministerie van Economische Zaken en ministerie van Infrastructuur en Milieu, Den Haag.

Rijkswaterstaat, 2019. Kader Ecologie en Cumulatie 3.0 t.b.v. uitrol van windenergie op zee 2030. Ministerie van Landbouw, Natuur en Voedselkwaliteit, Den Haag.

- Rodgers, J. A. and Schwikert, S. T. (2002) Buffer-zone distances to protect foraging and loafing waterbirds from disturbance by personal watercraft and outboard-powered boats. *Conserv. Biol.* 16: 216–224.
- Rydell, J., L. Bach, M.-J. Bubour-Savage, M. Green, L. Rodrigues, & A. Hedenström, 2010a. Bat mortality at wind turbines in Northwestern Europe. *Acta Chiropterologica* 12: 261–274.
- Rydell, J., L. Bach, M.-J. Bubour-Savage, M. Green, L. Rodrigues, & A. Hedenström, 2010b. Mortality of bats at wind turbines links to nocturnal insect migration? *European Journal of Wildlife Research* 56: 823-827.
- Rydell, J., L. Bach, P. Bach, L. Guia Diaz, J. Furmarkiewicz, N. Hagner-Wahlsten, E.-M. Kyheröinen, T. Lilley, M. Masing, M. Max Meyer , G. Pétersons, J. Šuba, V. Vasko, V. Vintulis, & A. Hedenström, 2014. Phenology of Migratory Bat Activity Across the Baltic Sea and the South-Eastern North Sea. *Acta Chiropterologica* 16(1): 139-147.
- Scharringa, C.J.G., W. Ruitenbeek & P.J. Zomerdijk, 2010. *Atlas van de Noord-Hollandse broedvogels 2005-2009. Samenwerkende Vogelwerkgroepen Noord-Holland (SVN) / Landschap Noord-Holland.*
- Schwemmer, P., Mendel, B., Sonntag, N., Dierschke, V. and Garthe, S. (2011) Effects of ship traffic on seabirds in offshore waters: implications for marine conservation and spatial planning. *Ecol. Appl.* 21: 1851–1860.
- Skov, H., S. Heinänen & M. Lazcny, 2015a. Offshore Wind Farm Eneco Luchterduinen Ecological monitoring of seabirds TConstr report. ENECO
- Skov, H., S. Heinänen, L. Nyborg & M. Lazcny, 2015b. Offshore Wind Farm Eneco Luchterduinen Ecological monitoring of seabirds T0 report. ENECO
- Skov, H., S. Heinänen, L., M. Lazcny & M. Chudzinska, 2017. Offshore Wind Farm Eneco Luchterduinen Ecological monitoring of seabirds T2 report. ENECO.
- Skov, H., S. Heinänen, T. Norman, R.M. Ward, S. Mendez-Roldan & I. Ellis, 2018. ORJIP Bird Collision and Avoidance Study. Final report–April 2018. The Carbon Trust, United Kingdom.
- Tasker, M.L., Jones, P.H., Dixon, T.J. & Blake, B.F. 1984 Counting seabirds at sea from ships: a review of methods employed and a suggestion for a standardized approach. *Auk* 101, 567-577.
- Thorpe, A. 2005. Recent Reports May to September 2005. *The Fulmar, Bulletin of the North Sea Bird Club* 109: 6.
- Van der Laar, F. 2007. Green light to birds: investigation into the effect of bird-friendly lighting. Report NAM location L15-FA-1. NAM, Assen, The Netherlands.
- Van der Wal, J.T., A. Gyimesi, R.C. Fijn & M. Scholl, 2015. 2nd Iteration: Effect of turbine capacity on collision numbers for three large gull species, based on revised density data, when

assessing cumulative effects of offshore wind farms on birds in the Southern North Sea.
Additional note to IMARES report number C166/14.

Vanermen N., E.W.M. Stienen, W. Courtens, T. Onkelinx, M. Van de Walle & H. Verstraete 2013. Bird monitoring at offshore wind farms in the Belgian part of the North Sea - Assessing seabird displacement effects. Rapport van het Instituut voor Natuur- en Bosonderzoek 2013 (INBO.R.2013.755887). INBO, Brussel.

Vanermen N., T. Onkelinx, W. Courtens, M. Van de Walle, H. Verstraete & E.W.M. Stienen, 2014. Seabird avoidance and attraction at an offshore wind farm in the Belgian part of the North Sea. *Hydrobiologia* doi: 10.1007/s10750-014-2088-x.

van der Laar, F. 2007. Green light to birds: investigation into the effect of bird-friendly lighting. Report NAM location L15-FA-1. NAM, Assen, The Netherlands.

van der Wal, J.T., A. Gyimesi, R.C. Fijn & M. Scholl, 2015. 2nd Iteration: Effect of turbine capacity on collision numbers for three large gull species, based on revised density data, when assessing cumulative effects of offshore wind farms on birds in the Southern North Sea.
Additional note to IMARES report number C166/14.

van der Wal, J.T., M.E.B. van Puijenbroek & M.F. Leopold, 2018. Cumulatieve effecten van offshore wind parken: habitatverlies zeevogels: update voor vijf zeevogelsoorten tot 2030 (No. C059/18). Wageningen Marine Research.

Walls, R., S. Canning, G. Lye, L. Givens, C. Garrett & J. Lancaster 2013. Analysis of Marine Environmental Monitoring Plan Data from the Robin Rigg Offshore Wind Farm, Scotland. (Operational Year 1). E.ON Climate & Renewables & Natural Power publication 1022038.

Winter, H.V., G. Aarts & O.A. van Keeken 2010. Residence time and behaviour of sole and cod in the Offshore Wind farm Egmond aan Zee (OWEZ). Report number
OWEZ_R_265_T1_20100916. Imares Wageningen UR.

Wood, K.A., J.L. Newth, G.M. Hilton, B.A. Nolet & E.C. Rees, 2016. Inter-annual variability and long-term trends in breeding success in a declining population of migratory swans. *Journal of avian biology* 47(5): 597-609.

<http://bd.eionet.europa.eu/article17/reports2012>

Hoofdstuk 7 Onderwaterleven (zie ook de literatuurlijst die bij bijlage 5 hoort)

- Ainslie, M.A. (2010). Principles of sonar performance modeling. Springer-Verlag Berlin Heidelberg. pp. 707.
- Aarts, G., J. Cremer, R. Kirkwoord, J.T. van der Wal, J. Matthiopoulos & S. Brasseur, 2016. Spatial distribution and habitat preference of harbour seal (*Phoca vitulina*) in the Dutch North Sea. Wageningen Marine Research report C118/16.
- Aarts, G., S. Brasseur, S. Geelhoed, R. van Bemmelen & M. Leopold, 2013. Grey and harbour seal spatiotemporal distribution along the Dutch West coast. IMARES report C103/13.
- ACRB, 2013. Mogelijke morfologische effecten van het Prinses Amaliawindpark. Rapport in opdracht van Prinses Amaliawindpark. ACRB.
- Andersson, M., Andersson, S., Ahlsén, J., Andersson, B., Hammar, J., Persson, L., Pihl, J., Sigray, P., Wikström, A., 2017. A Framework for Regulating Underwater Noise During Pile Driving. Report by Vindval. pp 115.
- Arts, F.A., S. Lilipaly & R.C.W. Strucker, 2014. Watervogels en zeezoogdieren in de Zoute Delta 2012/2013. RWS Centrale Informatievoorziening BM 14.11.
- ASCOBANS, 2011. Summary Record of the 18th Meeting of the Advisory Committee. UN Campus, Bonn, Germany, 4-6 May 2011.
- de Backer, A. en Hostens K. 2018. Soft sediment epibenthos and fish monitoring at the belgian offshore wind farm area:situation 6 and 7 years after constructiion. In: Degræer, S., Brabant, R., Rumes, B. & Vigin, L. (eds). 2018. *Environmental Impacts of Offshore Wind Farms in the Belgian Part of the North Sea: Assessing and Managing Effect Spheres of Influence*. Brussels: Royal Belgian Institute of Natural Sciences, OD Natural Environment, Marine Ecology and Management, 136 p.
- Bergman M., G. Duineveld, R. Daan, M. Mulder & S. Ubels, 2012. Impact of OWEZ wind farm on the local macrobenthos community macrobenthos. Final report OWEZ_R_261_T2_20121010. NoordzeeWind – Wageningen Imares.
- Bergman M., G. Duineveld, R. Daan, M. Mulder & S. Ubels, 2012. Impact of OWEZ wind farm on the local macrobenthos community macrobenthos. Final report OWEZ_R_261_T2_20121010. NoordzeeWind – Wageningen Imares.
- BMU, 2013. Bundesministerium für Umwelt, Naturschutz und Reaktorsicherheit. Konzept für den Schutz der Schweinswale vor Schallbelastungen bei der Errichtung von Offshore-Windparks in der deutschen Nordsee (Schallschutzkonzept).
- Bochert R and Zettler ML 2006. Effect of electromagnetic fields on marine organisms, in Offshore Wind Energy. Springer. ;223-234.

Bolle LJ, de Jong CAF, Bierman SM, van Beek PJG, van Keeken OA, Wessels PW et al. (2012) Common Sole Larvae Survive High Levels of Pile-Driving Sound in Controlled Exposure Experiments. *PLoS ONE* 7(3): e33052. doi:10.1371/journal.pone.0033052

Bolle, L.J., E. Blom, C.A.F. de Jong, M.B. Halvorsen, R. Hoek , C.J.G van Damme, P.W. Wessels, H.V. Winter, C.M. Woodley, C. Hinrichs, T. Huijer, M. Lohman, I. Pennock, 2014. Sub-lethal effects of pile-driving sounds on juvenile sea bass. IMARES WageningenUR report number C111.14, IJmuiden, Nederland.

Bolle, L.J., C. A. F. de Jong, S. M. Bierman, P .J. C. van Beck, P. W. Wessels, E. Blom, C .J. G. van Damme, H. V. Winter & R. P. A. Dekeling, 2016. Effect of pile-driving sounds on the survival of larval fish. *Advances in Experimental Medicine and Biology* 875: 91 - 100.

Bos O.G., R. Witbaard, M. Lavaleye, G. van Moorsel, L.R. Teal, R. van Hal, T. van der Hammen, R. ter Hofstede, R. van Bemmelen, R.H. Witte, S. Geelhoed & E.M. Dijkman, 2011. Biodiversity hotspots on the Dutch Continental Shelf. IMARES Wageningen UR. Report number C071/11.

Bos O.G., A. Gittenberger, I. de Boois, M. van Asch, J.T van der Wal, J. Cremer, B. van der Hoorn, S. Pieterse, P.A.J. Bakker, 2016. Soortenlijst Nederlandse Noordzee. Wageningen Marine Research rapport C125/16. DOI: <https://doi.org/10.18174/401117>

Bouma S., & W. Lengkeek 2009. Development of underwater flora- and fauna communities on hard substrates of the offshore wind farm Egmond aan Zee (OWEZ). Report OWEZ_R_266_T1_20090126

Bouma S., & W. Lengkeek 2009. Development of underwater flora- and fauna communities on hard substrates of the offshore wind farm Egmond aan Zee (OWEZ). Report OWEZ_R_266_T1_20090126.

Bouma S. & W. Lengkeek 2012. Benthic communities on hard substrates of the offshore wind farm Egmond aan Zee (OWEZ). Including results of samples collected in scour holes. Report OWEZ_R_266_T1_20120206_hard_substrate.

Brandt, M.J., A-C. Dragon, A. Diederichs, M.A. Bellmann, V. Wahl, W. Piper, J. Nabe-Nielsen & G. Nehls, 2018. Disturbance of harbour porpoises during construction of the first seven offshore wind farms in Germany. *Mar. Ecol. Prog. Ser.* 596: 213 – 232.

Brasseur, S.M.J.M., M. Scheidat, G.M. Aarts, J.S.M. Cremer & O.G. Bos, 2008. Distribution of marine mammals in the North Sea for the generic appropriate assessment of future offshore wind parks. IMARES report C046/08.

Brasseur, S., T. van Polanen Petel, G. Aarts, E. Meesters, E. Dijkman & P. Reijnders, 2010. Grey seals (*Halichoerus grypus*) in the Dutch North sea: population ecology and effects of wind farms. IMARES report No C137/10.

Brasseur, S. Aarts, G. Meersters, E. Polanen Petel, t. van, Dijkman, E. Cremer, J. & Reijnders, P. Habitat preferences of harbour seals in the Dutch coastal area: analysis and estimate of effects of offshore wind farms (2012). Rapport: OWEZ R 252 T1 20120130, 30-01-2012.

Camphuysen, C.J., 2011. Lesser Black-backed Gulls nesting at Texel. Foraging distribution, diet, survival, recruitment and breeding biology of birds carrying advanced GPS loggers. NIOZ-Report 2011-05. Royal Netherlands Institute for Sea Research, Texel

Camphuysen, C.J. & M.L. Siemensma, 2011. Conservation plan for the Harbour Porpoise *Phocoena phocoena* in The Netherlands: towards a favourable conservation status. NIOZ Report 2011-07, Royal Netherlands Institute for Sea Research, Texel.

Carpenter, J.R., Merckelbach L., Callies U., Clark S., Gaslikova L., Baschek B., 2016. Potential Impacts of Offshore Wind Farms on North Sea Stratification. PLOS One 11(8) e0160830.

Casper, B.M., Popper, A.N., Matthews, F., Carlson, T.J., Halvorsen, M.B. 2012. Recovery of Barotrauma Injuries in Chinook Salmon from Exposure to Pile Driving Sound. PLoS ONE 7: e39593

Casper BM, Halvorsen MB, Matthews F, Carlson TJ, Popper AN 2013. Recovery of Barotrauma Injuries Resulting from Exposure to Pile Driving Sound in Two Sizes of Hybrid Striped Bass. PLoS ONE 8: e73844. doi:10.1371/journal.pone.0073844 De Robertis et al., 2003

Casper, B.M., Halvorsen, M.B., Carlson, T.J. and Popper, A.N., 2017. Onset of barotrauma injuries related to number of pile driving strike exposures in hybrid striped bass. The Journal of the Acoustical Society of America, 141(6), pp.4380-4387.

CBS, PBL & Wageningen UR, 2012. Ruimtelijke verdeling biodiversiteit in de Noordzee, 1991-2010 (indicator 2159, versie 01 , 25 september 2012) .
<http://www.compendiumvoordeleefomgeving.nl/> . CBS; Planbureau voor de Leefomgeving en Wageningen UR, Den Haag, Bilthoven en Wageningen.

Coates, D.A., Deschutter, Y., Vincx, M. and Vanaverbeke, J., 2014. Enrichment and shifts in macrobenthic assemblages in an offshore wind farm area in the Belgian part of the North Sea. Marine environmental research, 95, pp.1-12.

Coates, D.A., Van Hoey, G., Colson, L., Vincx, M. and Vanaverbeke, J., 2015. Rapid macrobenthic recovery after dredging activities in an offshore wind farm in the Belgian part of the North Sea. *Hydrobiologia*, 756(1), pp.3-18.

Compendium voor de leefomgeving, www.clo.nl, 2016, 2018 en 2019.

Coolen, J.W., Bos, O.G., Glorius, S., Lengkeek, W., Cuperus, J., van der Weide, B. and Agüera, A., 2015. Reefs, sand and reef-like sand: A comparison of the benthic biodiversity of habitats in the Dutch Borkum Reef Grounds. *Journal of Sea Research*, 103, pp.84-92.
 Coolen, J.W.P., 2017. *North Sea reefs: benthic biodiversity of artificial and rocky reefs in the southern North Sea* (Doctoral dissertation, Wageningen University).

Couperus, B., E. Winter, O. van Keeken, T. van Kooten, S. Tribuhl & D. Burggraaf, 2010. Use of high resolution sonar for near_turbine fish observations (DIDSON) – We@Sea 2007_002.

IJmuiden, IMARES

Costa, D.P., 2012. A bioenergetics approach to developing a population consequences of acoustic disturbance model. In: Popper AN, Hawkins A (eds) "The effects of noise on aquatic life. Advances in experimental medicine and biology." Springer Science and Business Media, New York, NY, p. 423–426.

Craeymeersch J.A., Witbaard R., Dijkman E., Meesters H.W.G., 2008. Ruimtelijke en temporele patronen in de diversiteit van macrobenthische infauna op het Nederlands Continentaal Plat. Report number C070/80 IMARES.

Daan N., 2005. Aanzet tot een systeembeschrijving Noordzee: de visgemeenschap. RIVO rapport C012/05.

Daan R., Mulder M & Bergman M.J.N., 2009. Impact of windfarm OWEZ on the local macrobenthos community. Report OWEZ_R_261_T1_20091216.

Dähne, M., J. Tougaard, J. Carstensen, A. Rose & J. Nabe-Nielsen, 2017. Bubble curtains attenuate noise from offshore wind farm construction and reduce temporary habitat loss for harbour porpoises. Mar Ecol Prog Ser 580: 221–237.

Damveld, J. H., van der Reijden, K. J., Cheng, C., Koop, L., Haaksma, L. R., Walsh, C. A. J., et al. (2018). Video transects reveal that tidal sand waves affect the spatial distribution of benthic organisms and sand ripples. Geophysical Research Letters, 45, 11,837–11,846.
<https://doi.org/10.1029/2018GL079858>.

De Robertis, A., C.H. Ryer, A. Veloza & R.D. Brodeur, 2003. Differential effects of turbidity on prey consumption of piscivorous and planktivorous fish. Canadian Journal of Fisheries and Aquatic Sciences 60(12): 1517-1526.

Debusschere, E., B. De Coensel, A. Bajek, D. Botteldooren, K. Hostens, J. Vanaverbeke, S. Vandendriessche, K. Van Ginderdeuren, M. Vincx & S. Degraer, 2014. In situ mortality experiments with juvenile sea bass (*Dicentrarchus labrax*) in relation to impulsive sound levels caused by pile driving of windmill foundations. PLoS one 9(10): e109280. Debusschere, E., Hostens, K., Adriaens, D., Ampe, B., Botteldooren, D., De Boeck, G., De Muynck, A., Sinha, A.K., Vandendriessche, S., Van Hoorebeke, L. and Vincx, M., 2016. Acoustic stress responses in juvenile sea bass *Dicentrarchus labrax* induced by offshore pile driving. Environmental Pollution, 208, pp.747-757.

Debusschere E, De Coensel B, Bajek A, Botteldooren D, Hostens K, Vanaverbeke J et al. (2014) In Situ Mortality Experiments with Juvenile Sea Bass (*Dicentrarchus labrax*) in Relation to Impulsive Sound Levels Caused by Pile Driving of Windmill Foundations. PLoS ONE 9(10): e109280. doi:10.1371/journal.pone.0109280

Debusschere, E., B. De Coensel, A. Bajek, D. Botteldooren, K. Hostens, J. Vanaverbeke, S. Vandendriessche, K. Van Ginderdeuren, M. Vincx & S. Degraer, 2014. In situ mortality experiments with juvenile sea bass (*Dicentrarchus labrax*) in relation to impulsive sound levels

caused by pile driving of windmill foundations. *PLoS one* 9(10): e109280. Debusschere, E., Hostens, K., Adriaens, D., Ampe, B., Botteldooren, D., De Boeck, G., De Muynck, A., Sinha, A.K., Vandendriessche, S., Van Hoorebeke, L. and Vincx, M., 2016. Acoustic stress responses in juvenile sea bass *Dicentrarchus labrax* induced by offshore pile driving. *Environmental Pollution*, 208, pp.747-757.

Derweduwen, J., Vandendriessche, S. and Hostens, K., 2016. Effects of Belgian wind farms on the epibenthos and fish of the soft sediment. Environmental impacts of offshore wind farms in the Belgian part of the North Sea: Environmental impact monitoring reloaded. Operationele Directie Natuurlijk Milieu.

Diederichs, A., H. Pehlke, G. Nehls, M. Bellmann, P. Gerke, J. Oldeland, C. Grunau, S. Witte & A. Rose, 2014. Entwicklung und Erprobung des Großen Blasenschleiers zur Minderung der Hydroschallemissionen bei Offshore-Rammarbeiten. BMU Förderkennzeichen 0325309A/B/C, BioConsult SH, Husum.

E-Connection project bv, (2008) Milieueffectrapport Offshore Windpark Q4-WP. Bunnik, oktober 2008.

Eigaard, O.R., F. Bastardie, M. Breen, G.E. Dinesen, N.T. Hintzen, P. Laffargue, L.O. Mortensen, J.R. Nielsen, H.C. Nilsson, F.G. O'Neill, H. Polet, D.G. Reid, A. Sala, M. Sköld, C. Smith, T.K. Sørensen, O. Tully, M. Zengin, A.D. Rijnsdorp 2016 Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design and dimensions. *ICES J Mar Sci*; 73 (suppl_1): i27-i43. doi: 10.1093/icesjms/fsv099

Emma, B., 2016. A Review of the Evidence of Electromagnetic Field (Emf) Effects on Marine Organisms. *Res. Rev. J Ecol. Environ. Sci.* 4(4):22-26).

Fey, D.P., Jakubowska, M., Greszkiewicz, M., Andrulewicz, E., Otremba, Z. and Urban-Malinga, B., 2019. Are magnetic and electromagnetic fields of anthropogenic origin potential threats to early life stages of fish?. *Aquatic Toxicology*, 209, pp.150-158.

Floeter, J., van Beusekom JE, Auch D, Callies U, Carpenter J, Dudeck T, Eberle S, Eckhardt A, Gloe D, Hänselmann K, Hufnagl M., 2017. Pelagic effects of offshore wind farm foundations in the stratified North Sea. *Progress in Oceanography*. 156:154-73.

Galatius A, S.M.J.M. Brasseur, P.J.H. Reijnders, T. Borchardt, U. Siebert, M. Stede, S. Ramdohr, L.F. Jensen & J. Teilmann (2012) Trilateral Seal Expert Group (TSEG). Aerial Surveys of Harbour Seals in the Wadden Sea in 2012. Substantial increases in moult counts. Common Wadden Sea Secretariate (CWSS).

Geelhoed, S., Scheidat, M., Aarts, G., van Bemmelen, R., Janinhoff, N., Verdaat, H. & Witte, R., 2011. Shortlist masterplan wind aerial surveys of harbour porpoises on the Dutch Continental Shelf. IMARES, Wageningen Report number C103/11.

Geelhoed, S.C.V., Scheidat, M., van Bemmelen, R.S.A. & Aarts, G., 2013. Abundance of harbour porpoises (*Phocoena phocoena*) on the Dutch Continental Shelf, aerial surveys in July

2010- March 2011. Lutra 56(1): 45-57.

Geelhoed, S.C.V., Scheidat, M., van Bemmelen, 2014, Marine mammal surveys in Dutch waters in 2013, Report number C027/14.

Geelhoed, S.C.V., S. Lagerveld & J.P. Verdaat, 2015. Marine mammal surveys in Dutch North Sea waters in 2015. IMARES report C189/15.

Geelhoed, S.C.V., N. Janinhoff, S. Lagerveld, L.S. Lehnert & J.P. Verdaat, 2018. Marine mammal surveys in Dutch North Sea waters in 2017. Wageningen Marine Research (University & Research centre), WMR report C030/18.

Goodson, A. D. and Sturtivant, C. R. 1996. Sonar characteristics of the harbour porpoise (*Phocoena phocoena*): source levels and spectrum. – ICES Journal of Marine Science, 53: 465–472.

Grontmij, 2008. Offshore Windpark Q10. Documentnummer 13/99088008/CvD.

Hal, R. van, B. Couperus, S. Fassler, S. Gastauer, B. Griffioen, N. Hintzen, L. Teal, O. van Keeken & E. Winter 2012. Monitoring- and Evaluation Program Near Shore Wind farm (MEP-NSW). Fish community. IMARES Report C059/12.

OWEZ_R_264_T1_20121215_final_report_fish. Imeres Wageningen UR.

Haelters J. & C.J. Camphuysen 2009. The Harbour Porpoise (*Phocoena phocoena* L.) in the southern North Sea: Abundance, threats, research- and management proposals. Royal Belgian Institute of Natural Sciences (RBINS), department Management Unit of the North Sea Mathematical Models (MUMM) & Royal Netherlands Institute for Sea Research (NIOZ) Project financed by IFAW (International Fund for Animal Welfare) - IFAW Internationaler Tierschutz-Fonds GmbH, IFAW, Brussels.

Haelters, J., Kerckhof, F., Jauniaux, T. & Degraer, S. (2012b). The Grey Seal (*Halichoerus grypus*) as a Predator of Harbour Porpoises (*Phocoena phocoena*)? Aquatic Mammals 38(4): 343-353.

Haelters, J., Van Roy, W., Vigin, L. & Degraer, S., 2012. The effect of pile driving on harbour porpoises in Belgian waters. In: S. Degraer, R. Brabant & B. Rumes (Eds.). Offshore windfarms in the Belgian part of the North Sea: heading for an understanding of environmental impacts. Royal Belgian Institute of Natural Sciences, Department MUMM, Chapter 9: 127-143.

Halvorsen, M.B., Casper, B.M., Woodley, C.M., Carlson, T.J., Popper, A.N., 2012a. Threshold for Onset of Injury in Chinook Salmon from Exposure to Impulsive Pile Driving Sounds. PLoS ONE 7: e38968

Halvorsen, M.B., Casper, B.M., Matthews, F., Carlson, T.J., Popper, A.N. 2012b. Effects of exposure to pile-driving sounds on the lake sturgeon, Nile tilapia and hogchoker. Proc. R. Soc. B 279: 4705-4714

Heinis F., C.J. de Jong & Werkgroep Onderwatergeluid, 2015. Cumulatieve effecten van impulsief onderwatergeluid op zeezoogdieren. Rapport TNO 2015 R10335.

Heinis, F., C.A.F. de Jong, S. von Benda-Beckmann & B. Binnerts, 2019. Framework for Assessing Ecological and Cumulative Effects – 2018; Cumulative effects of offshore wind farm construction on harbour porpoises. HWE rapport: 18.153RWS_KEC2018, January 2019

Holtmann, S.E., A. Groenewold, K.H.M. Schrader, J. Asjes, J.A. Craeymeersch, G.C.A. Duineveld, A.J. van Bostelen en J. van der Meer, 1996. Atlas of the zoobenthos of the Dutch Continental Shelf, Ministry of transport, Public Works and Water Management, North Sea Directorate, Rijswijk, pp 244.

Jansen, O.E. (2013) Fishing for food: Feeding ecology of harbour porpoises *Phocoena phocoena* and white-beaked dolphins *Lagenorhynchus albirostris* in Dutch waters. PhD thesis, Wageningen University, Wageningen, NL

Jansen, H.W. & C.A.F. de Jong 2014. Underwater noise measurements in the North Sea in and near the Princess Amalia Wind Farm in operation. TNO report - TNO 2013 R11916. Den Haag, the Netherlands.

Jarvis S., Allen J., Proctor N., Crossfield A., Dawes O., Leighton A., McNeill L., Musk W., 2004. Dutch Wind Farms: NSW Lot 1 Benthic Fauna. Final Report 19 February 2004. Institute of Estuarine & Coastal Studies (IECS) The University of Hull. Reference No: ZBB607.2-F-2004.

JASCO Research Ltd (2006). Vancouver Island Transmission Reinforcement Project: Atmospheric and Underwater Acoustics Assessment. Report prepared for British Columbia Transmission Corporation 49 pp., <http://www.eao.gov.bc.ca>

Jaspers Faijer, M. & P. van Dam, 2013. Cumulative Effects of Offshore Wind Farms in the North Sea. – Discussion Report. Project number 712030. Pondera Consult, Hengelo.

De Jong, M.F., M.J. Baptist, H.J. Lindeboom & P. Hoekstra, 2015. Relationships between macrozoobenthos and habitat characteristics in an intensively used area of the Dutch coastal zone. ICES Journal of Marine Science: Journal du Conseil 72(8): 2409-2422. De Robertis, A., C.H. Ryer, A. Veloza, R.D. Brodeur, 2003. Differential effects of turbidity on prey consumption of piscivorous and planktivorous fish. Canadian Journal of Fisheries and Aquatic Sciences 60: 1517-1526

De Jong, C & B. Binnerts, 2016. Onderwatergeluidberekeningen Net op Zee Hollandse Kust (zuid). Bijlage 2 bij Passende Beoordeling Net op zee Hollandse Kust (zuid). Passende beoordeling Net op zee Hollandse Kust (zuid). A&W-rapport 2263.

De Jong, M.F., M.J. Baptist, H.J. Lindeboom & P. Hoekstra, 2015. Relationships between macrozoobenthos and habitat characteristics in an intensively used area of the Dutch coastal zone. ICES Journal of Marine Science: Journal du Conseil 72(8): 2409-2422. De Robertis, A., C.H. Ryer, A. Veloza, R.D. Brodeur, 2003. Differential effects of turbidity on prey consumption of piscivorous and planktivorous fish. Canadian Journal of Fisheries and Aquatic Sciences 60: 1517-1526

Jongbloed, R.H., D.M.E. Slijkerman, R. Witbaard & M.S.S. Lavaleye, 2013. Ontwikkeling zeebodemintegriteit op het Friese Front en de Centrale Oestergronden in relatie tot bodemberoerende visserij: Verslag expert workshop. IMARES

Kamermans, P. & M. Dedert, 2012. Effect of variations in concentration of algae and silt on filtration and growth of the razor clam (*Ensis directus*, Conrad). IMARES Report C017/12. IMARES, Wageningen.

Kastelein, R.A., N. Steen, R. Gransier & C.A.F. de Jong, 2013. Brief Behavioral Response Threshold Level of a Harbor Porpoise (*Phocoena phocoena*) to an Impulsive Sound. *Aquatic Mammals* 39: 315–323.

Kastelein, R.A., Jennings, N., Kommeren, A., Helder-Hoek, L. and Schop, J., 2017. Acoustic dose-behavioral response relationship in sea bass (*Dicentrarchus labrax*) exposed to playbacks of pile driving sounds. *Marine Environmental Research*, 130, pp.315-324.

Kerckhof, F., De Mesel, I., Degraer, S. 2016. Do wind farms favour introduced hard substrata species. In: Degraer, S., Brabant, R., Rumes, B., Vigin, L. (Eds.) (2016). Environmental impacts of offshore wind farms in the Belgian part of the North Sea: Environmental impact monitoring reloaded. Royal Belgian Institute of Natural Sciences, OD Natural Environment, Marine Ecology and Management Section. 287 pp.

Kirschvink , J.L.(1990) Geomagnetic sensitivity in cetaceans; An update with live stranding records in the United States. In J.A. Thomas & R.A. Kastelein, sensory abilities of cetaceans; laboratory and field evidence (Vol. 196, pp. 639 – 649). New York: Plenum.

Krone, R., Dederer, G., Kanstinger, P., Krämer, P., Schneider, C. and Schmalenbach, I., 2017. Mobile demersal megafauna at common offshore wind turbine foundations in the German Bight (North Sea) two years after deployment-increased production rate of Cancer pagurus. *Marine environmental research*, 123, pp.53-61.

Leewis L. & Klink A. 2017. Prinses Amalia Windturbine park 2017. Statistical comparison of benthic fauna inside and outside the Prinses Amalia Wind Park; a preliminary analysis. Korte notitie in opdracht van Rijkswaterstaat. Eurofins AquaSense.

Leewis, L. & Klink A., 2017. Prinses Amalia Windturbine park 2017. Statistical comparison of benthic fauna inside and outside the Prinses Amalia Wind Park; a preliminary analysis. Korte notitie in opdracht van Rijkswaterstaat. Eurofins AquaSense.

Leewis, L., Klink, A.D., Verduin, E.C. (2018) Benthic development in and around offshore wind farm Prinses Amalia Wind Park near the Dutch coastal zone be-fore and after construction (2003-2017) A statistical analysis (Reference RWS: 4500264484), Rijkswaterstaat, 65 pp.

Lefaible, N., Braeckman, U., Moens, T., 2018 Effects of wind turbine foundations on surrounding macrobenthic communities. In: Degraer, S., Brabant, R., Rumes, B. & Vigin, L. (eds). 2018. *Environmental Impacts of Offshore Wind Farms in the Belgian Part of the North*

Sea: Assessing and Managing Effect Spheres of Influence. Brussels: Royal Belgian Institute of Natural Sciences, OD Natural Environment, Marine Ecology and Management, 136 p.

Lengkeek, W., K. Didderen, M. Dorenbosch, S. Bouma, H. W. Waardenburg, 2013. Biodiversiteit van kunstmatige substraten. Een inventarisatie van 10 scheepswrakken op het NCP. Rapport 13-226, Bureau Waardenburg, Culemborg

Leonhard, S.B. & Pedersen, J. 2006. Benthic communities at Horns Rev before, during and after Construction of Horns Rev Offshore Wind Farm Vattenfall. Report number: Final Report/Annual Report 2005: 134.

Leopold, M.F., M.J. Baptist, L. IJsseldijk & B. Engels Rapport C096/13
Lindeboom, H., J. Geurts van Kessel & L. Berkenbosch, 2005. Gebieden met bijzondere ecologische waarden op het Nederlands Continentaal Plat. Rapport RIKZ/2005.008.

Lindeboom, H.J., Kouwenhoven, H.J., Bergman, M.J.N., Bouma, S., Brasseur, S., Daan, R., Fijn, R.C., de Haan, D., Dirksen, S., van Hal, R., Hille Ris Lambers, R., ter Hofstede, R., Krijgsveld, K.L., Leopold M. & Scheidat M., 2011. Short-term ecological effects of an offshore wind farm in the Dutch coastal zone; a compilation. Environmental Research Letters 6 035101 doi: 10.1088/1748-9326/6/3/035101.

Lock K., Faasse M. and Vanagt T.J., 2014. Assessment of the soft sediment fauna sixyears after construction of the Princess Amalia wind farm. eCOAST report 2013002.

Love, M.S., Nishimoto, M.M., Clark, S. and Bull, A.S., 2015. Identical response of caged rock crabs (*Genera Metacarcinus* and *Cancer*) to energized and unenergized undersea power cables in Southern California, USA. *Bulletin, Southern California Academy of Sciences*, 114(1), pp.33-42.

Love, M.S., Nishimoto, M.M., Clark, S., McCrea, M. and Bull, A.S., 2017. Assessing potential impacts of energized submarine power cables on crab harvests. *Continental Shelf Research*, 151, pp.23-29.

Meager, J.J., T. Solbakken, A.C. Utne-Palm, T. Oen, 2005. Effects of turbidity on the reactive distance, search time, and foraging success of juvenile Atlantic cod (*Gadus morhua*). Canadian Journal of Fisheries and Aquatic Sciences 62: 1978-1984

Nabe-Nielsen, J. R.M. Sibly, J. Tougaard, J. Teilmann & S. Sveegaard, 2014. Effects of noise and by-catch on a Danish harbour porpoise population. *Ecol. Modell.* 272, 242–251.

Nedelec, S.L., Campbell, J., Radford, A.N., Simpson, S.D. and Merchant, N.D., 2016. Particle motion: the missing link in underwater acoustic ecology. *Methods in Ecology and Evolution*, 7(7), pp.836-842.

New, L.F., J. S. Clark, D. P. Costa, E. Fleishman, M. A. Hindell, T. Klanjšček, D. Lusseau, S. Kraus, C. R. McMahon, P. W. Robinson, R. S. Schick, L. K. Schwarz, S. E. Simmons, L.

Thomas, P. Tyack, J. Harwood. 2014. Using short-term measures of behaviour to estimate long-term fitness of southern elephant seals. MEPS 496:99-108.

Nielsen, N.H., J. Teilmann, S. Svegaard, R.G. Hansen, M-H.S. Sinding, R. Dietz & M.P. Heide-Jørgensen, 2018. Oceanic movements, site fidelity and deep diving in harbour porpoises from Greenland show limited similarities to animals from the North Sea. Mar. Ecol. Prog. Ser. 597, 259 – 272

NMFS – National Marine Fisheries Service, 2016. Technical Guidance for Assessing the Effects of Anthropogenic Sound on Marine Mammal Hearing: Underwater Acoustic Thresholds for Onset of Permanent and Temporary Threshold Shifts. NOAA Technical Memorandum NMFS-OPR-55.

Öhman, M.C., Sigray, P. and Westerberg, H., 2007. Offshore windmills and the effects of electromagnetic fields on fish. *AMBIOS: A journal of the Human Environment*, 36(8), pp.630-634.

Platteeuw M., N.F. van der Ham & J.E. den Ouden, 1994. Zeetrekellingen in Nederland in de jaren tachtig. *Sula* 8(1/2, special issue): 1-203.

Popper, A.N., M.E. Smith, P.A. Cott, B.W. Hanna, A.O. MacGillivray, M.E. Austin & D.A. Mann, 2005. Effects of exposure to seismic airgun use on hearing of three fish species. *The Journal of the Acoustical Society of America* 117(6): 3958-3971.

Popper, A. N., and Hawkins, A. D. (eds) (2014). "The effects of noise on aquatic life II," Springer Science+Business Media, LLC, New York.

Popper, A. N., and Hawkins, A. D. (eds), 2014. "The effects of noise on aquatic life II," Springer Science+Business Media, LLC, New York.

Reiss, H., S.P.R. Greenstreet, K. Sieben, S. Ehrich, G.J. Piet, F. Quirijns, L. Robinson, W.J. Wolff, I. Kröncke, 2009. Effects of fishing disturbance on benthic communities and secondary production within an intensively fished area. *Marine Ecology Progress Series* , 394: 201–213

Reubens J., 2011. Windmolens in de Noordzee: een kraamkamer voor de schaarse kabeljauw. *Energieneuws* 27 juli 2012.

Reubens, J.T., Degraer, S. and Vincx, M., 2014. The ecology of benthopelagic fishes at offshore wind farms: a synthesis of 4 years of research. *Hydrobiologia*, 727(1), pp.121-136.

Rijnsdorp, A. D. (Designer), & Bos, O. G. (Designer). (2017). Beoordeling van het effect van bodemvisserij. Digital or Visual Products, Wageningen Marine Research. Retrieved from <http://edepot.wur.nl/429295>

Roach, M., Cohen, M., Forster, R., Revill, A.S., Johnson, M. (ed. S. Degraer), 2018. The effects of temporary exclusion of activity due to wind farm construction on a lobster (*Homarus gammarus*) fishery suggests a potential management approach. *ICES Journal of Marine Science*, 75(4), pp.1416-1426

- Roberts, L., Cheesman, S., Breithaupt, T. and Elliott, M., 2015. Sensitivity of the mussel *Mytilus edulis* to substrate borne vibration in relation to anthropogenically generated noise. *Marine Ecology Progress Series*, 538, pp.185-195.
- Roberts, L., Harding, H.R., Voellmy, I., Bruintjes, R., Simpson, S.D., Radford, A.N., Breithaupt, T. and Elliott, M., 2016, July. Exposure of benthic invertebrates to sediment vibration: from laboratory experiments to outdoor simulated pile-driving. In *Proceedings of Meetings on Acoustics* 4ENAL (Vol. 27, No. 1, p. 010029). ASA.
- Roberts, L. and M. Elliott, 2017. Good or bad vibrations? Impacts of anthropogenic vibration on the marine epibenthos. *Science of the Total Environment*, 595, pp.255-268.
- Rosen, D.A.S., A.J. Winship & L.A. Hoopes, 2007. Thermal and digestive constraints of foraging behaviour in marine mammals. *Philosophical Transactions of the Royal Society B: Biological Sciences* 362: 2151-2168.
- Scheidat M, Verdaat H, Aarts G (2012) Using aerial surveys to estimate density and distribution of harbour porpoises in Dutch waters. *Journal of Sea Research* 69:1-7.
- Scheidat, M., R. Leaper, M. van den Heuvel-Greve & A. Winship, 2013. Setting Maximum Mortality Limits for Harbour Porpoises in Dutch Waters to Achieve Conservation Objectives. *Open Journal of Marine Science* 2013, 3.
- Scott, K., Harsanyi, P. & A.R. Lyndon, 2018. Understanding the effects of electromagnetic field emissions from Marine Renewable Energy Devices (MREDs) on the commercially important edible crab, *Cancer pagurus* (L.). *Marine pollution bulletin*, 131, pp.580-588.
- Sierra-Flores, R., Atack, T., Migaud, H. and Davie, A., 2015. Stress response to anthropogenic noise in Atlantic cod *Gadus morhua* L. *Aquacultural engineering*, 67, pp.67-76.
- Slavik, K., Lemmen C., Zhang W., Kerimoglu O., Klingbeil K., Wirtz K.W. 2017. The large scale impact of o_shore windfarm structures on pelagic primary production in the southern North Sea. arXiv preprint arXiv:1709.02386
- Snoek, R., R de Swart, K. Didderen, W. Lengkeek, M. Teunis, 2016. Potential effects of electromagnetic fields in the Dutch North Sea, Phase 1: Desk study.
- Southall, B.L., A.E. Bowles, W.T. Ellison, J.J. Finneran, R.L. Gentry, C.R. Greene Jr., D. Kastak, D.R. Ketten, J.H. Miller, P.E. Nachtigall, W.J. Richardson, J.A. Thomas & P.L. Tyack, 2007. Marine Mammal Noise Exposure Criteria: Initial Scientific Recommendations. *Aquatic Mammals*, 33(4), 411–521.
- Sparling, C.E. & M.A. Fedak, 2004. Metabolic rates of captive grey seals during voluntary diving. *J Exp Biol* 207: 1615-1624.

Sparling, C.E., J-Y. Georges, S.L. Gallon, M. Fedak & D. Thompson, 2007. How long does a dive last? Foraging decisions by breath-hold divers in a patchy environment: a test of a simple model. *Animal Behaviour* 74: 207-218.

Spiga, I., Caldwell, G.S. and Bruintjes, R., 2016, July. Influence of Pile Driving on the Clearance Rate of the Blue Mussel, *Mytilus edulis* (L.). In Proceedings of Meetings on Acoustics 4ENAL (Vol. 27, No. 1, p. 040005). ASA.

Spiga, I., Aldred, N. and Caldwell, G.S., 2017. Anthropogenic noise compromises the anti-predator behaviour of the European seabass, *Dicentrarchus labrax* (L.). *Marine Pollution Bulletin*, 122(1-2), pp.297-305.

Sveegaard, S., J. Teilmann, P. Berggren, K.N. Mouritsen, D. Gillespie & J. Tougaard, 2011. Acoustic surveys confirm the high-density areas of harbor porpoises found by satellite tracking. *ICES Journal of Marine Science* 68: 929–936.

Taormina, B., Bald, J., Want, A., Thouzeau, G., Lejart, M., Desroy, N. & A. Carlier, 2018. A review of potential impacts of submarine power cables on the marine environment: Knowledge gaps, recommendations and future directions. *Renewable and Sustainable Energy Reviews*, 96, pp.380-391.

Thomsen, F., K. Lüdemann, R. Kafemann & W. Piper, 2006. Effects of offshore wind farm noise on marine mammals and fish. Biola, Hamburg, Germany on behalf of COWRIE Ltd 62.

Thomson 2015. MaRVEN – Environmental Impacts of Noise, Vibrations and Electromagnetic Emissions from Marine Renewable Energy.

Tougaard, J., A.J. Wright & P.T. Madsen, 2015. Cetacean noise criteria revisited in the light of proposed exposure limits for harbour porpoises. *Marine Pollution Bulletin* 90: 196–208.

Tricas, T. & A. Gill, 2011. Effects of EMFs from undersea power cables on elasmobranchs and other marine species. Normandeau Associates, Inc report. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Regulation, and Enforcement, Pacific OCS Region, Camarillo, CA. OCS Study BOEMRE 2011-09.

Troost, K., Perdon, K., Van Zwol, J., Jol, J. & Van Asch M., 2017. Schelpdierenbestanden in de Nederlandse kustzone in 2017. CVO rapport: 17.014.

Van Damme, C.J.G., Hoek, R., Beare, D., Bolle, L.J., Bakker, C., van Barneveld, E., Lohman, M., Os-Koomen, E., Nijssen, P., Pennock I.,& Tribuhl, S. 2011. Shortlist Master plan Wind Monitoring fish eggs and larvae in the Southern North Sea: Final report Part A. Report number C098/11 IMARES.

Van Denderen, P.D., T. van Kooten & A.D. Rijnsdorp, 2013. When does fishing lead to more fish? Community consequences of bottom trawl fisheries in demersal food webs. *Proceedings of the Royal Society of London B: Biological Sciences* 280(1769): 20131883.

Van Denderen P. D. Hintzen N. T. Rijnsdorp A. D. Ruardij P. van Kooten T. 2014. Habitat-specific effects of fishing disturbance on Benthic species richness in marine soft sediments. *Ecosystems*, 17: 1216–1226.

Van Denderen, P.D., Bolam, S.G., Hiddink, J.G., Jennings, S., Kenny, A., Rijnsdorp, A.D. and Van Kooten, T., 2015. Similar effects of bottom trawling and natural disturbance on composition and function of benthic communities across habitats. *Marine Ecology Progress Series*, 541, pp.31-43.

Van Deurs, M., T.M. Grome, M. Kaspersen, H. Jensen, C. Stenberg, T.K. Sørensen, J. Støtrup, T. Warnar & H. Mosegaard, 2012. Short-and long-term effects of an offshore wind farm on three species of sandeel and their sand habitat. *Marine Ecology Progress Series* 458: 169-180.

van der Reijden, K.J., Hintzen, N.T., Govers, L.L., Rijnsdorp, A.D. & H. Olff, 2018. North Sea demersal fisheries prefer specific benthic habitats. *PloS one*, 13(12), p.e0208338.

van Dijk, T.A., van Dalfsen, J.A., Van Lancker, V., van Overmeeren, R.A., van Heteren, S. & P.J. Doornenbal, 2012. Benthic habitat variations over tidal ridges, North Sea, the Netherlands. In *Seafloor geomorphology as benthic habitat* (pp. 241-249). Elsevier.

Van Duin C.F., Jaspers C.J., Kreft M., 2011. Offshore Windpark Q10. Milieueffectrapport. Grontmij, 2011.

van Duren, L.A., Gittenberger, A., Smaal, A.C., Koningsveld, M. van, Osinga, R., Cado van der Lelij, J.A., Vries, M.B. de., 2016. Rijke riffen in de Noordzee : verkenning naar het stimuleren van natuurlijke riffen en gebruik van kunstmatig hard substraat. Rapport 1221293-000, Deltares (2016)

Van Hal R., B. Couperus, S. Fassler, S. Gastauer, B. Griffioen, N. Hintzen, L. Teal, O. van Keeken, E. Winter, 2012. Monitoring- and Evaluation Program Near Shore Wind farm (MEP-NSW) - Fish community. IMARES Report C059/12
OWEZ_R_264_T1_20121215_final_report_fish.

Van Hal R., 2013. Round Fish Monitoring Princess Amalia Wind Farm. IMARES report number C117/13A.

Van Hal R., 2014. Demersal Fish Monitoring Princess Amalia Wind Farm. IMARES report number C125/14.

Van Hal, R., Griffioen, A.B. and van Keeken, O.A., 2017. Changes in fish communities on a small spatial scale, an effect of increased habitat complexity by an offshore wind farm. *Marine Environmental Research*, 126, pp.26-36.

Van Moorsel, G.W.N.M., 2003. Ecologie van de Klaverbank, Biota Survey 2002. Ecosub, Doorn, 154p.

Van Scheppingen Y. & A. Groenewold, 1990. De ruimtelijke verspreiding van het benthos in de

zuidelijke Noordzee. De Nederlandse kustzone overzicht 1988-1989. Rijkswaterstaat Directie Noordzee/Dienst getijdenwateren. MILZON-BENTHOS rapport 90-03.

Vanagt T., Van de Moortel L., Heusinkveld J., Faasse M., Pérez-Domínguez R. and Lock K., 2013. Assessment of the soft sediment fauna five years after construction of the Princess Amalia wind farm. eCOAST report 2012011-2.

Vanagt *et al.* 2014

Vanagt T. and Faasse M., 2014. Development of hard substratum fauna in the Princess Amalia Wind Farm. Monitoring six years after construction. eCOAST report 2013009.

Vanaverbeke, J., U. Braeckman, E. de Borger, N. Mavraki, E. Toussaint & H. Voet, 2018. Interactions between biological and physical effects modify local biogeographic cycling processes in offshore wind farms. Poster EMBS Ostende 2018

Vandendriessche, S., Derweduwen, J. and Hostens, K., 2015. Equivocal effects of offshore wind farms in Belgium on soft substrate epibenthos and fish assemblages. Hydrobiologia, 756(1), pp.19-35.

Westerberg, H., I. Lagenfelt, & H. Svedang. 2007. Silver eel migration behaviour in the Baltic. ICES Journal of Marine Science 64: 1457-1462.

Westerberg, H. & I. Lagenfelt, 2008. Sub-sea power cables and the migration behaviour of the European eel. Fisheries Management and Ecology. 15(5-6): p. 369-375.

Winter, H.V., G. Aarts & O.A. van Keeken 2010. Residence time and behaviour of sole and cod in the Offshore Wind farm Egmond aan Zee (OWEZ). Report number OWEZ_R_265_T1_20100916. Imares Wageningen UR.

Witbaard, R. & P. Kamermans, 2009. De bruikbaarheid van de klepstandmonitor op *Ensis directus* ten behoeve van de monitoring van aan zand winning gerelateerde effecten. NIOZ rapport 2009-10. NIOZ, Texel.

Wright, S.R., Lynam, C.P., Righton, D.A., Metcalfe, J., Hunter, E., Riley, A., Garcia, L., Posen, P. & Hyder, K., 2018. Structure in a sea of sand: fish abundance in relation to man-made structures in the North Sea. *ICES Journal of Marine Science*.

Zabar, Z., Birenbaum, L., Cheo, B.R., Joshi, P.N. & Spagnolo, A., 1992. A detector to identify a de-energized feeder among a group of live ones. IEEE transactions on power delivery, 7(4), pp.1820-1824. DOI: 10.1109/61.156984

Hoofdstuk 8 Scheepvaartveiligheid

Barentse, J. Nadere toelichting: Gevolgen van aanvaringen door de windturbine-installatie.
Jacobs Comprimo Nederland, juli 2000.

Ministerie van Infrastructuur en Milieu, Beleidsnota Noordzee 2016-2021, 2015

J.T.M. van Doorn, A.M. Duursma, Y.Koldenhof, J. Valstar, WIND OP ZEE 2030: Gevolgen voor scheepvaartveiligheid en mogelijke mitigerende maatregelen. MARIN, 31132-3-MSCN-rev.1.0, 13 mei 2019

Koldenhof, Y. & C. van der Tak. Risico vervoer (milieu)gevaarlijke stoffen op zee. MARIN, 19287.630/4, juli 2004.

Koldenhof, Y. & Kaufmman, K. Effecten op Scheepvaartveiligheid voor Windenergiegebied Hollandse Kust (west). MARIN, 31909-1-MSCN, 12 Juli 2019.

Rijkswaterstaat, Memo: Advies extra inpassingsmaatregelen scheepvaart voor de Routekaart windenergie op zee 2030 benodigt bovenop bestaand beleid en beheer, 17 mei 2019

Verdrag inzake de Internationale Bepalingen ter voorkoming van aanvaringen op zee (COLREGs). Londen, 1972.

Hoofdstuk 9 Landschap

Bishop, Ian. 2002. Determination of thresholds of visual impact: The case of wind turbines. Environment and Planning B: Planning and Design. 29. 707-718. 10.1068/b12854.

KNMI, 2012. Uurgegevens van het weer in Nederland. Gedownloade gegevens. Via <https://www.knmi.nl/nederland-nu/klimatologie/uurgegevens>.

Kyba, Christopher C. M.; Mohar, Andrej; Posch, Thomas (2017). "How bright is moonlight?". Astronomy & Geophysics. 58 (1): 1.31–1.32. doi:10.1093/astrogeo/atx025.

Lörzing, .H, Harbers, A., Breedijk, M., 2007. De zichtbaarheid van de Bellen van Zuylen-toren. Ruimtelijk Planbureau, Den Haag.

Martinez-Condo & Macknick, 2007: Venster op de Geest

Nierman et al, 2010, Beleving en Maatschappelijke aspecten zichtbaarheid windturbines Noordzee.

Pondera Consult, 2018, MER Kavel V en VI Windenergiegebied Hollandse Kust (noord) (2018), bijlage 10 – Zichtbaarheidsanalyse

Pondera Consult, 2018, MER Kavel V en VI Windenergiegebied Hollandse Kust (noord) (2018), bijlage 10 – Zichtbaarheidsanalyse

Pondera Consult, 2015, MER Kavels I t/m IV, Hollandse Kust (zuid) (2015)

Shang, Haidong & Bishop, Ian. (2000). Visual Thresholds for Detection, Recognition and Visual Impact in Landscape Settings. Journal of Environmental Psychology. 20. 125-140. 10.1006/jenvp.1999.0153.

Hoofdstuk 10 Overige gebruiksfuncties

Agentschap Telecom, 2017, Toetsingscriterium Straalverbindingen en Windturbines.

Besluit luchtverkeer, 2014, Geraadpleegd via <https://wetten.overheid.nl/BWBR0035899/2014-12-12>.

Verordening EU nr. 923/2012, UITVOERINGSVERORDENING (EU) Nr. 923/2012 VAN DE COMMISSIE van 26 september 2012

Brown, C. Offshore Wind Farm Helicopter Search and Rescue Trials Undertaken at the North Hoyle Wind Farm; Report of helicopter SAR trials undertaken with Royal Air Force 'C' Flight 22 Squadron on March 22nd 2005. Maritime and Coastguard Agency, 2005.

Decisio, 2016, Regionale effecten windmolenparken op zee, Maatschappelijke effecten en analyse regionaal economische impact

Ecorys, 2019, Kansen, risico's en kosten voor de visserij bij toestaan sleepnetvisserij in windenergiegebieden Eindrapportage, Rotterdam.

European Parliament, 2019, Conservation of fishery resources and protection of marine ecosystems through technical measures, Strasbourg, 16 April 2019

Green Giraffe, 2019, LCOE impact of seabed fishing in OWFs, 22 July 2019

Howard, M. & C. Brown. Results of the electromagnetic investigations and assessments of marine radar, communications and positioning systems undertaken at the North Hoyle wind farm by QinetiQ and the Maritime and Coastguard Agency, 2004.

IALA, Recommendation O-139 – The Marking of Man-made Offshore Structures December 2008.

Ministerie van Binnenlandse Zaken en Koninkrijksrelaties, 2019, Ontwerp Nationale Omgevingsvisie. Geraadpleegd via <https://www.denationaleomgevingsvisie.nl/actueel/artikelen+en+blogs/1381493.aspx>

Ministerie van Infrastructuur en Milieu, 2015, Uitwerking besluit doorvaart en medegebruik van windparken op zee in het kader van Nationaal Waterplan 2016 – 2021, december 2015

Mosselzaadinvanginstallaties. Geraadpleegd via <https://pdokviewer.pdok.nl/#> op 6 mei 2019.

NLOG, 2019. Geraadpleegd via <https://www.nlog.nl/kaart-boringen>, update april 2019

Noorzeeloket, 2018 Indicatieve routes recreatievaart Noordzee.

Radio Holland, Onderzoek naar radarverstoring door Prinses Amaliawindpark en Offshore Windpark Egmond aan Zee, Resultaten van de veldexperimenten in 2010 bij kalme zee, 2012

Howard et all., 2004, Results of the electromagnetic investigations and assessments of marine radar, communications and positioning systems undertaken at the North Hoyle wind farm by QinetiQ and the Maritime and Coastguard Agency, 15 November 2004.

Periplus Archeomare, 2018 Archaeological Desk Study Hollandse Kust (west) Wind Farm Zone, december 2018

Periplus Archeomare, 2019, Hollandse Kust (west) - An archeological assessment of geophysical survey results, juli 2019

Primo Marine, 2019, TRA-1 Consequences of possible sea-bed fishery in future offshore wind farms, 27 July 2019.

PwC, 2018, De economische bijdrage van windenergie op zee, Rapport voor het Ministerie van Economische Zaken en Klimaat. PricewaterhouseCoopers.

REAS Euro, 2018, UXO desk study: site data Hollandse Kust (west) wind farm zone.

Rijksoverheid, Nieuwsbericht 13-12-2011: Delen Noordzee verboden voor visserij door akkoord natuurbeweging, vissers en rijksoverheid. Geraadpleegd via:
<https://www.rijksoverheid.nl/actueel/nieuws/2011/12/13/delen-van-noordzee-verboden-voor-visserij-door-akkoord-natuurbeweging-vissers-en-rijksoverheid>

Spaven consulting. Wind turbines and radar: operational experience and mitigation measures. Report to a consortium of wind energy companies, december 2001.

Stichting de Noordzee, 2018, Gebiedsbescherming op de Nederlandse Noordzee – De stand van zaken in relatie tot de visserij, Januari 2018.

To70, 2017, Helicopter accessibility of oil & gas platforms near the offshore wind farms Hollandse Kust Zuid and Hollandse Kust Noord, The Hague.

To70, 2018, Helicopter accessibility ' Hollandse Kust (west)', 'IJmuiden Ver' and 'Ten noorden van de Waddeneilanden' Results.

van der Reijden et all, 2018, North Sea demersal fisheries prefer specific benthic habitats. PLoS ONE 13(12): e0208338. <https://doi.org/10.1371/journal.pone.0208338>.
<https://www.wur.nl/nl/nieuws/Noordzeevissers-tonen-voorkleur-voor-zeldzame-habitats.htm>.

Wageningen Marine Research, 2017, Memo Pilot Mosselteelt op de Noordzee, 12 juni 2017

Wageningen Economic Research, 2019 Wind op Zee: bepaling van de waarde van geplande windparkgebieden voor de visserij. Mol, Arie; Oostenbrugge, Hans van; Röckmann, Christine; Hintzen, Niels

WUR, 2010, Vangst visserij per km² in de omgeving van windenergiegebied Hollandse Kust (west) (peildatum 2007) Bron: LEI, Wageningen UR 2010

WUR, 2019, Geschiktheid zeewindparken voor maricultuur en passieve visserij - Een kwalitatieve beoordeling van geschiktheid van windparklocaties voor voedselproductie, Lisanne van den Bogaart; Marnix Poelman, juni 2019, WuR

Wageningen University, 2018,
<http://www.agrimatie.nl/PublicatiePage.aspx?subpubID=2526&themalID=2286&indicatorID=2880§orID=2860>, bron geraadpleegd in april 2019.

Hoofdstuk 11 Elektriciteitsopbrengst

Centraal Bureau voor de Statistiek, 2016. Hernieuwbare energie in Nederland 2015. Den Haag/Heerlen, 2016.

CBS, 2018a, Rendementen en CO₂ emissie elektriciteitsproductie 2016. URL: www.cbs.nl/nl-nl/achtergrond/2018/04/rendementen-en-co2-emissie-elektriciteitsproductie-2016

CBS, 2018b, Emissies van luchtverontreinigde stoffen volgens NEC richtlijnen

CBS, 2018c, Hernieuwbare energie in Nederland 2017, 2018

Chaouki Ghenai, 2012, Life Cycle Analysis of Wind Turbine, Sustainable Development - Energy, Engineering and Technologies - Manufacturing and Environment, Prof. Chaouki Ghenai (Ed.), ISBN: 978-953-51-0165-9, InTech, Available from:
<http://www.intechopen.com/books/sustainable-development-energy-engineering-andtechnologies-manufacturing-and-environment/life-cycle-analysis-of-wind-turbine>

Das Grüne Emissionshaus, augustus 2003

Karl R. Haapala and Preedanood Prempreeda, 2014, Comparative life cycle assessment of 2.0 MW wind Turbines. In: Int. J. Sustainable Manufacturing, Vol. 3, No. 2, 2014,

<http://www.energy.siemens.com/hq/en/renewable-energy/wind-power/epd.htm>

http://www.noordzeewind.nl/wp-content/uploads/2012/02/OWEZ_R_000_20101112_Operations_2009.pdf

<http://www.prinsesamaliawindpark.eu/nl/windpark.asp>

<https://www.eneco.nl/over-ons/projecten/windpark-luchterduinen/>

Hoofdstuk 12 Afweging

-